



11-177549

- 1 -

(Translation)

[Name of the Document] SPECIFICATION

[Title of the Invention] COMPOSITION FOR PROMOTING PASSIVE
EXTENSION OF BLADDER SMOOTH MUSCLE

[Claims]

[Claim 1] A composition for promoting passive extension of bladder smooth muscle, comprising adrenomedullin.

[Claim 2] A composition according to claim 1, used to ameliorate a urination disorder.

[Claim 3] A composition according to claim 2, wherein the urination disorder is a urinary incontinence selected from the group consisting of urge incontinence, reflex incontinence, and overflow incontinence.

[Claim 4] A composition according to claim 1, wherein the adrenomedullin is:

(a) a peptide comprising an amino acid sequence from Ser in position 13 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING; or

(b) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (a), and having an action of promoting extension of bladder smooth muscle.

[Claim 5] A composition according to claim 4, wherein the adrenomedullin is:

(c) a peptide comprising an amino acid sequence from Tyr in position 1 to Tyr in position 52 of SEQ ID NO: 2 in

RECEIVED
FEB 12 2004
TECH. CENTER 1600/2900

11-177549

- 2 -

SEQUENCE LISTING; or

(d) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (c), and having an action of promoting extension of bladder smooth muscle.

[Claim 6] A composition according to claim 5, wherein the adrenomedullin is:

(e) a peptide comprising an amino acid sequence from Ala in position -73 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING; or

(f) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (e), and having an action of promoting extension of bladder smooth muscle.

[Claim 7] A composition according to claim 6, wherein the adrenomedullin is:

(g) a peptide comprising an amino acid sequence from Met in position -94 to Leu in position 91 of SEQ ID NO: 2 in SEQUENCE LISTING; or

(h) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (g), and having an action of promoting extension of bladder smooth muscle.

[Claim 8] A composition according to any of claims 1 and 4 to 7, wherein the C-terminus of the adrenomedullin is amidated.

[Claim 9] A composition according to any of claims 1 and 4 to 7, wherein Gly is added to the C-terminus of the adrenomedullin.

11-177549

- 3 -

[Claim 10] A composition according to any of claims 1 and 4 to 7, wherein in the adrenomedullin, Cys in position 16 and Cys in position 21 of SEQ ID NO: 2 in SEQUENCE LISTING are crosslinked.

[Claim 11] A composition according to claim 10, wherein the crosslink is a disulfide bond.

[Claim 12] A composition according to claim 10, wherein the crosslink is a $-\text{CH}_2-\text{CH}_2-$ bond.

[Detailed Description of the Invention]

[0001]

[Field of the Invention]

The present invention relates to a composition for promoting extension of smooth muscle of the urinary bladder, comprising adrenomedullin.

[0002]

[Prior Art]

Urinary incontinence is a common but very severe condition which mostly causes patients to be embarrassed, encounter difficulties, and be driven to despair. Clearly, there is a strong demand for a reliable and safe method of treating urinary incontinence. To date such a demand has not been satisfied to an appropriate level.

[0003]

Urinary incontinence refers to a condition in which urine involuntarily flows out during the storage phase, and which is caused when there is a functional or organic abnormality in one or both of the urinary bladder and the urethra.

11-177549

- 4 -

Urinary incontinence occurs when bladder smooth muscle is involuntarily contracted so that the internal pressure of the urinary bladder is increased, or when urethral closure pressure created by the urethral sphincter and a supporting tissue surrounding the urethra is too weak to repel the internal pressure of the normal urinary bladder. Urinary incontinence is divided into several types depending on the pathology. Broad types are: urge incontinence; reflex incontinence; overflow incontinence (poorly compliant bladder), stress incontinence, total incontinence; and nocturnal enuresis.

[0004]

Urge incontinence is a condition in which urine involuntarily flows out accompanying a strong urge to urinate, or a symptom that, after feeling an urge to urinate, a patient cannot resist urine outflow and wets before reaching a toilet. These are divided into a motor type and a sensory type. Motor urge incontinence is caused by a disorder of an inhibitory pathway for the micturition reflex, or excitation of an excitatory pathway, a representative example of which is neurogenic bladder due to a lesion, such as for example a cerebrovascular disorder and a brain tumor. Representative examples of sensory urge incontinence include cystitis and urethritis.

[0005]

Reflex incontinence is a condition in which when the urinary bladder is filled with urine to some extent without a normal urge to urinate, the urinary bladder is reflectively contracted, so that urine involuntarily flows out. Reflex incontinence includes neurogenic bladder due to injury of the spinal cord above the urination center in the sacral spinal cord, urinary incontinence of infants, and the like.

11-177549

- 5 -

[0006]

Overflow incontinence is a condition in which since urine cannot be sufficiently excreted, the urinary bladder is excessively filled with urine and the urine gradually flows out. Overflow incontinence includes neurogenic bladder (poorly compliant bladder) due to injury of peripheral nerves, and disorders of the passage of the lower urinary tract due to prostatic hypertrophy or cancer.

[0007]

Stress incontinence is a symptom in which when a patient strains in sneezing or coughing, laughs, runs, or the like, so that abdominal pressure is rapidly increased, urine flows out without contraction of the urinary bladder. The increase in the abdominal pressure leads to a raise in the internal pressure of the urinary bladder. In this case, if it exceeds the urethral closure pressure, urine flows out. Females more often suffer from this disorder. A major cause is that supporting tissues surrounding the urethra are seriously weakened by parturition or aging, so that urethral closure pressure cannot be sufficiently generated.

[0008]

Total incontinence is a condition characterized by dysfunction of the urethral sphincter, and in which urine flows out from the urethra at all times irrespective of the presence or absence of abdominal pressure. The cause is injury of the urethral sphincter caused by trauma of the pelvis or surgery of the prostate.

[0009]

Nocturnal enuresis is also called bed-wetting, which

11-177549

- 6 -

is a condition in which patients of 4 or more years old, which is the age at which the habit of urinating is established, unconsciously void urine during sleep though they do not have an organic abnormality in the urinary tract or the nerve system and can urinate normally without urinary incontinence on awakening. This is caused by the premature inhibitory mechanism of the central nerve system for the micturition reflex.

[0010]

At present, anticholinergic agents are generally used for treatment of the disease showing urine storage disorders: (1) neurogenic bladder patients who have urinary incontinence due to the hyperactivity of urinary bladder and involuntarily urinate; (2) neurogenic bladder patients who do not have abnormal urinary bladder contraction but have a poorly compliant urinary bladder in which the internal pressure of the urinary bladder gradually increased as urine is filled; (3) one subset of patients who chiefly complain of pollakiuria; etc. However, clinically, despite the efficacy against urine storage disorders, when the anticholinergic agents are actually used, urinary bladder contraction is inhibited in urination so that urination disorders are often exacerbated to cause problems of side effects, such as increased residual urine and anuresis.

[0011]

As described above, clearly, urinary incontinence is one of today's major diseases, but current therapeutic methods are not satisfactory. There is a demand for a novel drug for treating urinary incontinence. The term "urination disorder" as used herein refers to an abnormal urination condition, such as urinary incontinence, caused by insufficient extension of

11-177549

- 7 -

bladder smooth muscle. Examples of urination disorders include urinary incontinence (e.g., urge incontinence), frequent urination, and nocturnal pollakiuria. An agent capable of promoting extension of bladder smooth muscle would be expected to help bladder smooth muscle extend during storage phase to reduce the internal pressure of the urinary bladder. Thus, such an agent would be considered to be useful as drugs for treatment of urinary incontinence and other symptoms relating to urination.

[0012]

It has been known that adrenomedullin has a vasodilatory action. For example, Nakamura et al., Jpn. J. Pharmacol. 67, 259-262 (1995) has reported in Figure 1 that contracted mesenteric artery is extended by addition of adrenomedullin in a concentration-dependent manner. However, vasodilation cannot be considered to be identical with passive extension of muscle of the urinary bladder. For example, Nishimura et al., British J. Pharmacology, 120, 193-200 (1997) describes in Figure 6 that addition of adrenomedullin to the urinary bladder does not cause contraction or extension (active extension) of the urinary bladder.

[0013]

[Problems to be Solved by the Invention]

The present invention is intended to solve the above-described problems. The objective of the present invention is to provide a novel agent for promoting passive extension of bladder smooth muscle.

[0014]

[Means for Solving the Problems]

The inventors found that adrenomedullin originally

11-177549

- 8 -

identified as a peptide having a hypotensive action does not directly extend bladder smooth muscle, but has an action of promoting passive extension of the urinary bladder wall due to urine storage, i.e., an action of promoting extension of bladder smooth muscle, and based on that finding, completed the present invention.

[0015]

Adrenomedullin does not inhibit contraction of the urinary bladder due to acetylcholine (i.e., urinary bladder contraction in urination) and therefore, can provide a therapeutic agent for ameliorating only a urine storage disorder without inhibiting urinary bladder contraction in urination and substantially without side effects.

[0016]

A composition of the present invention for promoting passive extension of bladder smooth muscle comprises adrenomedullin. The composition of the present invention may be used to ameliorate a urination disorder. The urination disorder may be a urinary incontinence selected from the group consisting of urge incontinence, reflex incontinence, and overflow incontinence.

[0017]

In one embodiment, the adrenomedullin may be any of the following peptides: (a) a peptide comprising an amino acid sequence from Ser in position 13 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING; (b) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (a), and having an action of promoting extension of bladder smooth muscle; (c) a peptide comprising an amino acid sequence from

11-177549

- 9 -

Tyr in position 1 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING; (d) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (c), and having an action of promoting extension of bladder smooth muscle; (e) a peptide comprising an amino acid sequence from Ala in position -73 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING; (f) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (e), and having an action of promoting extension of bladder smooth muscle; (g) a peptide comprising an amino acid sequence from Met in position -94 to Leu in position 91 of SEQ ID NO: 2 in SEQUENCE LISTING; and (h) a peptide consisting of an amino acid sequence having one or several amino acid deleted, substituted, or added in the amino acid sequence (g), and having an action of promoting extension of bladder smooth muscle.

[0018]

In another embodiment, the C-terminus of the adrenomedullin may be amidated, or Gly may be added thereto.

[0019]

In another embodiment, in the adrenomedullin, Cys in position 16 and Cys in position 21 of SEQ ID NO: 2 in SEQUENCE LISTING may be crosslinked. The crosslink may be a disulfide bond or a $-CH_2-CH_2-$ bond.

[0020]

[Embodiments of the Invention]

When carrying out the present invention, protein separation and analysis methods, recombinant DNA techniques, and assays, which are known in the art, may be employed unless

11-177549

- 10 -

otherwise specified.

[0021]

I. Definition

Hereinafter, the terms used herein to explain the present invention will be described.

[0022]

An adrenomedullin is a peptide originally isolated from human pheochromocytoma as a peptide having a hypotensive action. The term "adrenomedullin" as used herein is not limited to the particular peptide, but includes peptides having substantial homology in the amino acid sequence with that peptide. Examples of the homologous peptides include species mutants and allelic mutants. Human-derived adrenomedullin comprises an amino acid sequence from Tyr in position 1 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING. (The peptide consisting of an amino acid sequence from Met in position -94 to Leu in position 91 of SEQ ID NO: 2 in SEQUENCE LISTING is believed to be preproadrenomedullin. The peptide obtained by processing of a signal peptide and consisting of an amino acid sequence from Ala in position -73 to Leu in position 91 of SEQ ID NO: 2 in SEQUENCE LISTING is believed to be proadrenomedullin. The peptide consisting of an amino acid sequence from Ser in position 13 to Tyr in position 52 of SEQ ID NO: 2 in SEQUENCE LISTING is an adrenomedullin fragment which has been confirmed to have a hypotensive action. Any of the above-described forms may be employed in the present invention.) Human-derived adrenomedullin may be encoded by a polynucleotide sequence from T in position 447 to C in position 602 of SEQ ID NO: 1 in SEQUENCE LISTING.

[0023]

11-177549

- 11 -

Porcine-derived adrenomedullin comprises an amino acid sequence from Tyr in position 1 to Tyr in position 52 of SEQ ID NO: 4 in SEQUENCE LISTING. Porcine-derived adrenomedullin may be encoded by a polynucleotide sequence from T in position 430 to C in position 585 of SEQ ID NO: 3 in SEQUENCE LISTING. Rat-derived adrenomedullin comprises an amino acid sequence from Tyr in position 1 to Tyr in position 50 of SEQ ID NO: 6 in SEQUENCE LISTING. Rat-derived adrenomedullin may be encoded by a polynucleotide sequence from T in position 433 to T in position 582 of SEQ ID NO: 5 in SEQUENCE LISTING.

[0024]

Clearly, human-derived peptides are preferable for human diseases or treatment of a human. However, homologous peptides derived from other mammals may also be employed for some purposes. Further, comparison with peptides derived from other mammals is important in obtaining a variant maintaining a desired activity of a human-derived peptide.

[0025]

Adrenomedullin used in the present invention is not necessarily limited to the above-described sequences, but includes, as subjects, homologous peptides having an amino acid sequence which has one or several amino acid deleted, substituted, or added in the above-described sequences and maintaining a desired activity.

[0026]

Amino acid conservative substitution is one preferable means for obtaining homologous peptides. Conservative substitution representatively includes substitutions conducted within the following groups: glycine,

11-177549

- 12 -

alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

[0027]

The homology between two amino acid sequences is determined by optionally introducing a gap to optimize residue matching. A peptide having an amino acid sequence, which has substantially homology with the amino acid sequence of human adrenomedullin, has representatively about 60% homology with the amino acid sequence of human adrenomedullin, preferably at least about 70%, more preferably at least about 80%, and in an especially preferable embodiment, at least about 90% or more. Software for determining homology is easily available.

[0028]

In the present invention, a peptide is by definition referred to "have an action of promoting extension of bladder smooth muscle" if the degree of extension of the urinary bladder is about 80% or more and preferably about 90% or more of the value indicated in the experimental sample of Example 1 described below when measured under substantially the same conditions as those of Example 1 below.

[0029]

The C-terminus of a peptide used in the present invention may or may not be amidated. "Amidation of C-terminus" refers to one of modification reactions of a peptide, in which the COOH group of the C-terminal amino acid of a peptide is changed to the form of CONH₂. A number of biologically active peptides functioning in vivo are first biosynthesized as a precursor protein having a larger molecular

11-177549

- 13 -

weight. The precursor protein is then matured by a modification reaction such as for example the amidation of the C-terminus during an intracellular transition process. The amidation is conducted by a C-terminal amidating enzyme acting on the precursor protein. The precursor protein always includes a Gly residue on the C-terminal side of a residue to be amidated, which is frequently followed by a basic amino acid sequence pair, such as for example Lys-Arg or Arg-Arg, on the C-terminal side (Mizuno, Seikagaku, Vol. 61, No. 12, pp. 1435-1461 (1989)).

[0030]

II. Adrenomedullin having an action of promoting extension of bladder smooth muscle

In the present invention, adrenomedullin is utilized as an effective component of a composition for promoting extension of bladder smooth muscle. Adrenomedullin may be isolated from naturally-occurring sources, produced using recombinant DNA techniques, or chemically synthesized.

[0031]

When adrenomedullin is isolated from naturally-occurring sources, purification may be conducted, for example, in the following way. For example, firstly human pheochromocytoma is pulverized to obtain a crude extract which is in turn subjected to various chromatography techniques for purification of adrenomedullin. In this case, by monitoring an increase in the cAMP activity of platelets, a fraction containing adrenomedullin of interest can be obtained. Method for isolation and purification of adrenomedullin are described in Japanese Laid-Open Publication No. 7-196693.

[0032]

11-177549

- 14 -

When adrenomedullin is produced using recombinant DNA techniques, the DNA sequence encoding a peptide of interest is expressed using various recombinant systems. Construction of expression vectors and preparation of transformants having appropriate DNA sequences are conducted by methods known in the art. Expression may be conducted using prokaryote systems or eukaryote systems.

[0033]

Prokaryote hosts used include *E. coli*, bacillus, and other bacteria. For such a prokaryote, plasmid vectors having replication sites and control sequences compatible with the hosts are used. For example, *E. coli* is typically transformed with a derivative of pBR322 which is a plasmid derived from *E. coli*. The control sequence herein is defined to include a promoter for initiation of transcription, an operator if necessary, and a ribosome binding site sequence. Such a control sequence includes generally used promoters such as for example β -lactamase and lactose promoter systems (Chang et al., *Nature* (1977) 198, 1056), tryptophan promoters (Goeddel et al., *Nucleic Acids Res.*, (1980) 8: 4057), and P_L promoters derived from λ and N-gene ribosome binding sites (Shimatake, *Nature* (1981) 292, 128).

[0034]

As a eukaryote host, yeast is used, for example. For such a eukaryote, a plasmid vector having a replication site and a control sequence compatible with the host is used. For example, yeast is transformed with pYEUra3 (Clontech). Other promoters useful in a yeast host include, for example, promoter classes for synthesizing a glycolytic enzyme, such as for example a promoter for 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* (1980) 255, 2073). Other promoters

- 15 -

11-177549

include those derived from an enolase gene or those derived from a Leu2 gene obtained from YEp13.

[0035]

Appropriate mammalian promoters include metallothionein, an early or late promoter derived from SV40, and other virus promoters such as for example promoters derived from polyoma virus, adenovirus II, bovine papilloma virus and avian sarcoma virus.

[0036]

A transformant can be obtained by introducing an expression vector into an appropriate host cell. A desired adrenomedullin can be obtained by culturing the transformant under appropriate conditions.

[0037]

Chemical synthesis of adrenomedullin may be conducted within a method known in the art. For example, adrenomedullin may be synthesized by a solid phase method using a peptide synthesizer. A C-terminal amidated peptide can be produced on a peptide synthesizer by condensing amino acids sequentially from the C-terminal amino acid to the N-terminal amino acid using a benzhydryl amine resin and a standard DCC/HOBt, and cutting out an intended peptide from the resultant peptide resin by a standard cleavage method (trifluoromethanesulfonic acid method).

[0038]

A C-terminal amidated adrenomedullin may be obtained by: chemically amidating a carboxyl group at the C-terminus of the peptide obtained by expression in a host; or preparing a peptide so as to have Gly added to the C-terminus of an intended

11-177549

- 16 -

amino acid sequence for amidation by an action of the above-mentioned C-terminal amidating enzyme.

[0039]

Alternatively, the peptide obtained by adding Gly to the C-terminus of adrenomedullin may be amidated due to an action of a C-terminal amidating enzyme in vivo as described above.

[0040]

A disulfide bond can be formed, for example, by oxidizing a peptide by air oxidization or with an appropriate oxidant. The substitution of the disulfide bond with a $-CH_2-CH_2-$ bond can be conducted by a well-known method (O. Keller et al., *Helv. Chim. Acta* (1974) 57: 1253). Generally, cleavage in the disulfide bond is avoided by substituting a $-CH_2-CH_2-$ bond for the disulfide bond, resulting in stabilization of the protein.

[0041]

Assay methods for action of promoting extension of bladder smooth muscle, which are known in the art, may be used to confirm that the thus-obtained adrenomedullin has an action of promoting extension of bladder smooth muscle. Examples of such assays include a method employing the urinary bladder isolated from any animal, and a method of measuring the internal pressure of the urinary bladder under anesthetization. When the urinary bladder isolated from a rat is used, for example, action of promoting extension of bladder smooth muscle may be assayed under the following conditions. The urinary bladder is isolated from a rat and cut into several pieces to obtain urinary bladder strips. While the resultant urinary bladder strips are immersed in a buffer solution, such as for

11-177549

- 17 -

example Tyrode's solution, extension or contraction of the urinary bladder is continuously examined using a measuring apparatus, such as for example an isometric transducer and an isotonic transducer. The urinary bladder is allowed to be extended in the presence or absence of a subject peptide, and extension of the urinary bladder is compared to judge whether or not the peptide has an action of promoting extension of bladder smooth muscle.

[0042]

III. Preparation of a composition for promoting extension of bladder smooth muscle

A composition of the present invention comprises an effective amount of adrenomedullin and may comprise any excipient known to those skilled in the art. Examples of the excipients include lactose, cornstarch, magnesium stearate, and alum.

[0043]

The composition of the present invention is prepared in accordance with methods known in the art.

[0044]

The composition of the present invention may be in any form. The composition of the present invention may be a solid, such as for example a tablet, a pill, a capsule, and a granule; or a liquid, such as for example an aqueous solution and a suspension. When the composition of the present invention is orally administered as a tablet, an excipient, such as for example lactose, cornstarch, and magnesium stearate, may be commonly used. When the composition of the present invention is orally administered as a capsule, an excipient, such as for example lactose and dried cornstarch, may be

11-177549

- 18 -

commonly used. In order to orally administer adrenomedullin as an aqueous suspension, the adrenomedullin may be used in combination with an emulsion or a suspension. The aqueous suspension may optionally contain a sweetner and an aroma chemical. When the composition of the present invention is intramuscularly, intraperitoneally, subcutaneously, or intravenously injected, adrenomedullin is dissolved in a sterilized solution to prepare a buffer solution which is in turn adjusted into an appropriate pH. When the composition of the present invention is intravenously administered, the composition is preferably isotonic.

[0045]

IV. Administration of a composition for promoting extension of bladder smooth muscle

The composition of the present invention may be administered in the form of a conventional peptide formulation as described in Remington's Pharmaceutical Sciences, Mack Publishing, Easton, PA. For example, the composition of the present invention may be administered orally, or alternatively parenterally, such as for example intravenous administration, intramuscular injection, intraperitoneal injection, and subcutaneous injection. The peptide may be administered by injection into the urinary bladder.

[0046]

When the composition of the present invention is administered into a human subject, typically, the dose per day can be appropriately determined by those skilled in the art by taking into consideration a patient's symptoms, severity, individual differences in sensitivity, weight, age, and the like. The composition of the present invention may be administered once a day or may be administered several times

11-177549

- 19 -

a day.

[0047]

[Examples]

Hereinafter, the action of adrenomedullin of the present invention as a drug for promoting extension of bladder smooth muscle will be more specifically described. The present invention is not limited to the following examples. Adrenomedullin used in the examples is a synthesized peptide consisting of an amino acid sequence from Tyr in position 1 to Tyr in position 50 of SEQ ID NO: 6 (available from Peptide Institute, Inc.).

[0048]

(Example 1: Effect of adrenomedullin on extension of the urinary bladder of a male rat)

8 to 16 weeks old male rats were sacrificed by hammering their heads. Thereafter, the rats were decapitated, followed by exsanguination. The urinary bladders were isolated from the rats. Each isolated urinary bladder was cut into four portions, thereby obtaining urinary bladder strips (Figure 1).

[0049]

The effect of adrenomedullin on the rat urinary bladder was examined by measuring contractions of the urinary bladder strips using an isotonic transducer TD-112S (manufactured by Nippon Kohden Corporation) where the tension was 1 g.

[0050]

The urinary bladder strips were firstly immersed in 30 ml of Tyrode's solution with 100 nM adrenomedullin (experimental sample) or without it (control sample). In this

11-177549

- 20 -

situation, the sections were attached to the isometric transducer where the tension was 1 g, to continuously measure relaxation of the urinary bladder. The composition of the Tyrode's solution is as follows: 139 mM NaCl, 2.7 mM KCl, 11.9 mM NaHCO₃, 2.6 mM MgCl₂·6H₂O, 0.4 mM NaH₂PO₄·2H₂O, 1.7 mM CaCl₂, and 5.5 mM glucose; pH 7.4).

[0051]

The above-described experiment was repeated five times. The results are shown in Figure 2. Plain circles indicate averages of the control sample, while solid colored circles indicate the results of the experimental sample. Vertical bars extending from the points indicate the standard deviation of a two-way analysis of variance. The vertical axis indicates the length of relaxation (mm), while the horizontal axis indicates time (minutes).

[0052]

As shown in Figure 2, when tension was applied to the urinary bladder in the presence of adrenomedullin to cause the urinary bladder to be extended, the result obtained is that the urinary bladder wall was more extended compared to the case of the absence of adrenomedullin. Normally, the urinary bladder is extended by urine filling therein during the urine storage phase, and the extension prevents the internal pressure of the urinary bladder from being increased, and remains it at a constant low value. A high expansibility of the urinary bladder is referred to as a high level of compliance. As a result of this example, it was demonstrated that adrenomedullin causes the urinary bladder to remain in a higher compliance state during the urine storage phase, thereby increasing the capacity of the urinary bladder.

11-177549

- 21 -

[0053]

(Example 2: Effect of adrenomedullin on static tension, and contraction due to acetylcholine of the urinary bladder of a male rat)

Urinary bladder strips were prepared in a manner similar to that of Example 1. The urinary bladder sections were attached to an isometric transducer FD pickup TB611T (manufactured by Nippon Kohden Corporation) in Tyrode's solution. Contraction of the urinary bladder was continuously measured. Firstly, 30 nM to 1 mM acetylcholine was added to the Tyrode's solution. As a result, contraction of the urinary bladder occurred (Figure 3). In Figure 3, the vertical axis indicates tension (unit: g), while the horizontal axis indicates time. Contraction induced by acetylcholine was confirmed, followed by washing out acetylcholine. Thereafter, 100 nM adrenomedullin was added to the Tyrode's solution. As a result, the urinary bladder was not contracted. Further, 30 nM to 1 mM acetylcholine was added to the Tyrode's solution. As a result, recontraction of the urinary bladder occurred. The contraction induced by acetylcholine alone before the addition of adrenomedullin was not significantly different from the contraction induced by acetylcholine in the presence of adrenomedullin.

[0054]

Therefore, the tested adrenomedullin did not affect the static tension of the urinary bladder, or exhibit an effect of preventing contraction due to acetylcholine.

[0055]

[Effect of the Invention]

According to the present invention, a composition for promoting extension of bladder smooth muscle, comprising

11-177549

- 22 -

adrenomedullin is provided. Such a composition is useful for ameliorating a urination disorder selected from the group consisting of urge incontinence, reflex incontinence, and overflow incontinence.

[0056]

[Sequence Listing]

SEQUENCE LISTING

<110> Shionogi & Co., Ltd

<120> Composition for promoting passive dilation of the smooth muscle at urinary bladder

<130> J199137130

<140>

<141>

<150> 6

<170> PatentIn Ver. 2.0

<210> 1

11-177549

- 23 -

<211> 1457

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (166).. (719)

<220>

<221> mat peptide

<222> (447).. (602)

<400> 1

ggcaccgagct ggalagaaca gctcaagcct lgccacttcg ggccttcac tgcagctggg 60

cttggacttc gzagttttgc cattgccaagt gggacgtcig agactttctc ctccaagtac 120

ttggcagatc actctcttag cagggtctgc gcttcgcagc cggg atg aag ctg gtt 176

Met Lys Leu Val

tcc gtc gcc ctg atg tac ctg ggt tcg ctc gcc ttc cta ggc gct gac 224

Ser Val Ala Leu Met Tyr Leu Gly Ser Leu Ala Phe Leu Gly Ala Asp

-90

-85

-80

-75

acc gct cgg tlg gat gtc gcg tcg gag ttt cga aag aag tgg aat aag 272

Thr Ala Arg Leu Asp Val Ala Ser Glu Phe Arg Lys Lys Trp Asn Lys

-70

-65

-60

tgg gct ctg agt cgt ggg aag agg gaa ctg cgg atg tcc agc agc tac 320

11-177549

- 24 -

Trp Ala Leu Ser Arg Gly Lys Arg Glu Leu Arg Met Ser Ser Ser Tyr

-56

-50

-45

ccc acc ggg ctc gct gac gtg aag gcc ggg cct gcc cag acc cti att 368

Pro Thr Gly Leu Ala Asp Val Lys Ala Gly Pro Ala Gln Thr Leu Ile

-40

-35

-30

cgg ccc cag gac atg aag ggt gcc tct cga agc ccc gaa gac agc agt 416

Arg Pro Gln Asp Met Lys Gly Ala Ser Arg Ser Pro Gln Asp Ser Ser

-25

-20

-15

ccg gat gcc gcc cgc atc cga gtc aag cgc tac cgc cag agc atg aac 464

Pro Asp Ala Ala Arg Ile Arg Val Lys Arg Tyr Arg Gln Ser Met Asn

-10

-5

-1 1

5

aac ttc cag ggc ctc cgg agc ttt ggc tgc cgc ttc ggg acg tgc acg 512

Asn Phe Gln Gly Leu Arg Ser Phe Gly Cys Arg Phe Gly Thr Cys Thr

10

15

20

gtg cag aag ctg gca cac cag atc tac cag ttc aca gat aag gac aag 560

Val Gln Lys Leu Ala His Gln Ile Tyr Gln Phe Thr Asp Lys Asp Lys

25

30

35

gac aac gtc gcc ccc agg agc aag atc agc ccc cag ggc tac ggc cgc 608

Asp Asn Val Ala Pro Arg Ser Lys Ile Ser Pro Gln Gly Tyr Gly Arg

40

45

50

cgg cgc cgg cgc tcc ctg ccc gag gcc ggc cgg ggt cgg act ctg gtg 656

Arg Arg Arg Arg Ser Leu Pro Glu Ala Gly Pro Gly Arg Thr Leu Val

- 25 -

11-177549

55

60

65

70

tct tct aag cca cca gca cac ggg gct cca gcc ccc ccc agt gga agt 704
 Ser Ser Lys Pro Gln Ala His Gly Ala Pro Ala Pro Pro Ser Gly Ser
 75 80 85

gct ccc cac ttt ctt taggattag ggcaccaagg tacaaggaa agtcgcgcaa 759
 Ala Pro His Phe Leu
 90

gcaccccgct agtcctccc gggacgaagg acttcccgag cggcgagggg accgggctct 819

gacagccclg cggagacctt ggttcggga ggcaccgicc ggcggcgagc tcgggcttg 879

caaggggccc tcttctggg agcttcgctt cttagcctt gctcaggigc aagtgcccca 939

ggggcgggg tgcagaagaa tccgagtgtt tgccaggctt aaggagagga gaaactgaga 999

aatgaatgct gagaccoccg gaggagggt ctgagccaca gccgtgctcg cccacaaact 1059

gatttctac ggcgtgicac cccaccaggc cgaagcctc actattactt gaatttcca 1119

aaacctaaag aggaaaagtg caatgcgtgt tglacatata gaggtaacta tcaatatita 1179

agttgttgc tgcaagatt tttttgtaa cticaaatat agagatatit ttgtacgita 1239

tatattgtat taaggacatt ttaasagcaa ttatatgtc ctccctatt ttaagacgig 1299

aatgltcag cagggtgtaa agttgttcgc cgcgtggaa gtgagtgtgt ttgtgtcat 1359

- 26 -

11-177549

gaaagagaaa gacigattac ciccigigig gaagaaggaa acaccgagtc tetgtataat 1419

ctattfacat aaaatgggig atatggaac agcaaacc

1457

<210> 2

<211> 185

<212> PRT

<213> Homo sapiens

<400> 2

Met Lys Leu Val Ser Val Ala Leu Met Tyr Leu Gly Ser Leu Ala Phe

-90

-85

-80

Leu Gly Ala Asp Thr Ala Arg Leu Asp Val Ala Ser Glu Phe Arg Lys

-75

-70

-65

Lys Trp Asn Lys Trp Ala Leu Ser Arg Gly Lys Arg Glu Leu Arg Met

-60

-55

-50

Ser Ser Ser Tyr Pro Thr Gly Leu Ala Asp Val Lys Ala Gly Pro Ala

-45

-40

-35

Gln Thr Leu Ile Arg Pro Gln Asp Met Lys Gly Ala Ser Arg Ser Pro

-30

-25

-20

-15

Glu Asp Ser Ser Pro Asp Ala Ala Arg Ile Arg Val Lys Arg Tyr Arg

-10

-5

-1 1

11-177549

- 27 -

Gln Ser Met Asn Asn Phe Gln Gly Leu Arg Ser Phe Gly Cys Arg Phe
5 10 15

Gly Thr Cys Thr Val Gln Lys Leu Ala His Gln Ile Tyr Gln Phe Thr
20 25 30

Asp Lys Asp Lys Asp Asn Val Ala Pro Arg Ser Lys Ile Ser Pro Gln
35 40 45 50

Gly Tyr Gly Arg Arg Arg Arg Arg Ser Leu Pro Glu Ala Gly Pro Gly
55 60 65

Arg Thr Leu Val Ser Ser Lys Pro Gln Ala His Gly Ala Pro Ala Pro
70 75 80

Pro Ser Gly Ser Ala Pro His Phe Leu
85 90

<210> 3

<211> 1493

<212> DNA

<213> Sus scrofa

<220>

<221> CDS

<222> (148).. (711)

- 28 -

11-177549

<220>

<221> mat peptide

<222> (430).. (585)

<400> 3

gcggaacagc tcgagccttg ccacctctag ttcttiacca cagcttggac gtccggggtt 60

tgccactgcc agaggagcgt ctacagcttc atcttcccaa atcttggcag atcaccctt 120

tagcagggtc tgcacatctc agccggg atg aag ctg gtt ccc gta gcc ctc atg 174

Met Lys Leu Val Pro Val Ala Leu Met

-90

tac ctg ggc tgc ctg gcc ttc ctg ggc gct gac aca gct cgg ctc gac 222

Tyr Leu Gly Ser Leu Ala Phe Leu Gly Ala Asp Thr Ala Arg Leu Asp

-85

-80

-75

-70

gtg gcg gca gag ttc cga aag aaa tgg aat aag tgg gct cta agt cgt 270

Val Ala Ala Glu Phe Arg Lys Lys Trp Asn Lys Trp Ala Leu Ser Arg

-65

-60

-55

gga aaa aga gaa ctt cgg ctg tcc agc agc tac ccc acc ggg atc gcc 318

Gly Lys Arg Glu Leu Arg Leu Ser Ser Ser Tyr Pro Thr Gly Ile Ala

-50

-45

-40

gac ttg aag gcc ggg cct gcc cag act gtc att cgg ccc cag gat gtg 366

Asp Leu Lys Ala Gly Pro Ala Gln Thr Val Ile Arg Pro Gln Asp Val

-35

-30

-25

- 29 -

11-177549

aag ggc tcc tct cgc agc ccc cag gcc agc att ccg gat gca gcc cgc 414

Lys Gly Ser Ser Arg Ser Pro Gln Ala Ser Ile Pro Asp Ala Ala Arg

-20

-15

-10

atc cga gtc aag cgc tac cgc cag agt atg aac aac ttc cag gcc ctg 462

Ile Arg Val Lys Arg Tyr Arg Gln Ser Met Asn Asn Phe Gln Gly Leu

-5

-1 1

5

10

cgg agc ttc ggc tgt cgc ttt ggg acg tgc acc gtg cag aag ctg gcc 510

Arg Ser Phe Gly Cys Arg Phe Gly Thr Cys Thr Val Gln Lys Leu Ala

15

20

25

cac cag atc tac cag ttc acg gac aaa gac aag gac gcc gtc gcc ccc 558

His Gln Ile Tyr Gln Phe Thr Asp Lys Asp Lys Asp Gly Val Ala Pro

30

35

40

cgg agc aag atc agc ccc cag gcc tac ggc cgc cgg cgc cga cgc tct 606

Arg Ser Lys Ile Ser Pro Gln Gly Tyr Gly Arg Arg Arg Arg Ser

45

50

55

ctg ccc gaa gcc agc ctg gcc cgg act ctg agg tcc cag gag cca cag 654

Leu Pro Glu Ala Ser Leu Gly Arg Thr Leu Arg Ser Gln Glu Pro Gln

60

65

70

75

gcc cac ggg gcc ccg gcc tcc ccg gcc cat caa gtg ctg gcc act ctg 702

Ala His Gly Ala Pro Ala Ser Pro Ala His Gln Val Leu Ala Thr Leu

80

85

90

ttt agg att taggcgcta ctgtggcagc agcgaacagt cgcgcattgca 751

11-177549

- 30 -

Phe Arg Ile

tcattgccggc gcttctctggg gcggggggct tcccgagacc gagccctca ggggctggg 811

cccgggcaga gacagcattg agagaccgag agtccgggag gcacagacca gcggcgagcc 871

ctgcattttc aggaaccctt cctgcttggg ggcagtgctt tcttcggctt aatccagccc 931

gggtcccccg gtgggggtg aggtgcaga ggaatccaa ggagtgctat ctgccaggct 991

cacggagagg agaaacgtg agtaaatgc itagacccc agggcagg gtctgagcca 1051

ctgcctgcc gcccaaac tgatttctga aggggataa cccaacagg gcgcaggctt 1111

cactattact tgaacttcc aaacctaga gaggaaaagt gcaatgtatg ttgtatataa 1171

agaggtaact atcaatattt agtttgttg ctgtcaagat tttttttgt aactcaaat 1231

atagagatat ttttgtactg tatatatgt attaagggca ttttaaaac attgtatgt 1291

tccctcccc tctattttaa tatgtgaatg tctcagcgag gtgtaacatt gtttcttgcg 1351

cgaaatgtga gagtgtgtgt gtgtgtgtgc gtgaaagaga gctaggatgc ctcttgggga 1411

agaagaaaac acctatctg tataatctat ttacataaaa tgggtgatat gcgaagtgc 1471

aaaccaataa actgtctcaa tg

1493

- 31 -

11-177549

<210> 4

<211> 188

<212> PRT

<218> Sus scrofa

<400> 4

Met Lys Leu Val Pro Val Ala Leu Met Tyr Leu Gly Ser Leu Ala Phe
 -90 -85 -80

Leu Gly Ala Asp Thr Ala Arg Leu Asp Val Ala Ala Glu Phe Arg Lys
 -75 -70 -65

Lys Trp Asn Lys Trp Ala Leu Ser Arg Gly Lys Arg Glu Leu Arg Leu
 -60 -55 -50

Ser Ser Ser Tyr Pro Thr Gly Ile Ala Asp Leu Lys Ala Gly Pro Ala
 -45 -40 -35

Gln Thr Val Ile Arg Pro Gln Asp Val Lys Gly Ser Ser Arg Ser Pro
 -30 -25 -20 -15

Gln Ala Ser Ile Pro Asp Ala Ala Arg Ile Arg Val Lys Arg Tyr Arg
 -10 -5 -1 1

Gln Ser Met Asn Asn Phe Gln Gly Leu Arg Ser Phe Gly Cys Arg Phe
 5 10 15

Gly Thr Cys Thr Val Gln Lys Leu Ala His Gln Ile Tyr Gln Phe Thr
 20 25 30

11-177549

- 32 -

Asp Lys Asp Lys Asp Gly Val Ala Pro Arg Ser Lys Ile Ser Pro Gln
35 40 45 50

Gly Tyr Gly Arg Arg Arg Arg Arg Ser Leu Pro Glu Ala Ser Leu Gly
55 60 65

Arg Thr Leu Arg Ser Gln Glu Pro Gln Ala His Gly Ala Pro Ala Ser
70 75 80

Pro Ala His Gln Val Leu Ala Thr Leu Phe Arg Ile
85 90

<210> 5

<211> 1376

<212> DNA

<213> Rattus norvegicus

<220>

<221> CDS

<222> (154).. (708)

<220>

<221> nat peptide

<222> (483).. (582)

<400> 5

tccagccctt accgctcccg gttctcgccg ttctcatcgc agtcagctct ggactttgcg 60

11-177549

- 33 -

ggttttgccg ctgtcagaag gacgtctcgg actttctcgt tcaagtgtt gacaactcac 120

ccattcagca gggtatcgga gcatcgctac aga atg aag ctg gtt tcc atc gcc 174

Met Lys Leu Val Ser Ile Ala

-90

ctg atg tta ttg ggt tcg ctg gcc gtt ctg ggc gcg gac acc gca cgg 222

Leu Met Leu Leu Gly Ser Leu Ala Val Leu Gly Ala Asp Thr Ala Arg

-85

-80

-75

ctc gac act tcc tgg cag ttc cga aag aag tgg aat aag tgg gcg cta 270

Leu Asp Thr Ser Ser Gln Phe Arg Lys Lys Trp Asn Lys Trp Ala Leu

-70

-85

-80

-55

agt cgt ggg aag agg gaa cta caa gcg tcc agc agc tac cct acg ggg 318

Ser Arg Gly Lys Arg Glu Leu Gln Ala Ser Ser Ser Tyr Pro Thr Gly

-50

-45

-40

ctc gtt gat gag aag aca gtc ccg acc cag act ctt ggg ctg cag gac 366

Leu Val Asp Glu Lys Thr Val Pro Thr Gln Thr Leu Gly Leu Gln Asp

-35

-30

-25

aag cag agc acg tct agc acc cca caa gcc agc act cag agc aca gcc 414

Lys Gln Ser Thr Ser Ser Thr Pro Gln Ala Ser Thr Gln Ser Thr Ala

-20

-15

-10

cac att cga gtc aaa cgc tac cgc cag agc atg aac cag ggg tcc cgc 462

His Ile Arg Val Lys Arg Tyr Arg Gln Ser Met Asn Gln Gly Ser Arg

11-177549

- 34 -

-5

-1 1

5

10

agc act gga tgc cgc ttt ggg acc tgc aca atg cag aaa ctg gct cac 510
 Ser Thr Gly Cys Arg Phe Gly Thr Cys Thr Met Gln Lys Leu Ala His
 15 20 25

cag atc tac cag ttt aca gac aaa gac aag gac ggc atg gcc ccc aga 558
 Gln Ile Tyr Gln Phe Thr Asp Lys Asp Lys Asp Gly Met Ala Pro Arg
 30 35 40

aac aag atc agc cct caa ggc tat ggc cgc cgg cgc cgg cgt tcc ctg 606
 Asn Lys Ile Ser Pro Gln Gly Tyr Gly Arg Arg Arg Arg Arg Ser Leu
 45 50 55

cca gag gtc ctg cga gcc cgg act gtg gag tcc tcc cag gag cag acc 654
 Pro Gln Val Leu Arg Ala Arg Thr Val Gln Ser Ser Gln Gln Gln Thr
 60 65 70

cac tca gct cca gcc tcc ccg gcg cac caa gac atc tcc aga gtc tct 702
 His Ser Ala Pro Ala Ser Pro Ala His Gln Asp Ile Ser Arg Val Ser
 75 80 85 90

agg tta taggtcggg tggcagcatt gaacagtcgg gcgagtaicc cattggcgcc 758
 Arg Leu

tgcggaaica gagagcttcg caccctgagc ggactgagac aatcttgag agatctgcct 818

ggctggccct aggggagcca gaggaaacca agatcaagcc aggtcacgt cagaaaccga 878

11-177549

- 35 -

gaattacagg ctgatactct ctccgggcag gggctcagc cactgccctg cccgcata 938

aactggtttt ctacggggc ataccgtca ttacttactt gaacttcca aaacctagcg 998

aggaaaagtg caatgctgt taccagcca aggttaacta tcatattta gttgttgt 1058

gtcaagaggt ttttttttt gtaacttcaa atatatagaa atattttgt acgttatata 1118

tgtatttaag ggcattttta agcatttata ttgtcactt cccctattt aagaagtga 1178

tgtctcagca aggtgtaagg ttgttgggt ccgtgtgtgt gtgtgtgtgt gtgtgtgtgt 1238

gtgtgtgtgt gtgtgtgtta ggtggagagc gccgtattac cggcgttga tgaagaaaa 1298

acattgtgtc ttctataatc tatttacata aaatagtga tcgtggaaaa agcaaaccac 1358

ttaacigtct caatgtgt 1376

<210> 6

<211> 186

<212> PRT

<213> Rattus norvegicus

<400> 6

Met Lys Leu Val Ser Ile Ala Leu Met Leu Leu Gly Ser Leu Ala Val

-90

-85

-80

Leu Gly Ala Asp Thr Ala Arg Leu Asp Thr Ser Ser Gln Phe Arg Lys

11-177549

- 36 -

-75

-70

-65

Lys Trp Asn Lys Trp Ala Leu Ser Arg Gly Lys Arg Glu Leu Gln Ala

-60

-55

-50

Ser Ser Ser Tyr Pro Thr Gly Leu Val Asp Glu Lys Thr Val Pro Thr

-45

-40

-35

-30

Gln Thr Leu Gly Leu Gln Asp Lys Gln Ser Thr Ser Ser Thr Pro Gln

-25

-20

-15

Ala Ser Thr Gln Ser Thr Ala His Ile Arg Val Lys Arg Tyr Arg Gln

-10

-5

-1 1

Ser Met Asn Gln Gly Ser Arg Ser Thr Gly Cys Arg Phe Gly Thr Cys

5

10

15

Thr Met Gln Lys Leu Ala His Gln Ile Tyr Gln Phe Thr Asp Lys Asp

20

25

30

35

Lys Asp Gly Met Ala Pro Arg Asn Lys Ile Ser Pro Gln Gly Tyr Gly

40

45

50

Arg Arg Arg Arg Arg Ser Leu Pro Glu Val Leu Arg Ala Arg Thr Val

55

60

65

Glu Ser Ser Gln Glu Gln Thr His Ser Ala Pro Ala Ser Pro Ala His

70

75

80

- 37 -

11-177549

Gln Asp Ile Ser Arg Val Ser Arg Leu

85

90

[Brief Description of the Drawings]

[Figure 1] Figure 1 is a schematic diagram showing a method for dissecting the urinary bladder. Ends of the urinary bladder were cut off along solid lines as shown in Figure 1(a), and the urinary bladder was cut open along dashed lines. Four urinary bladder sections were obtained from the cut-open urinary bladder along three solid lines as shown in Figure 1(b).

[Figure 2] A graph showing the results of measurement of the effect of adrenomedullin on extension of the urinary bladder where the applied tension is 1 g. White circles indicate controls, while black circles indicate the case of addition of adrenomedullin.

[Figure 3] A graph showing the results of measurement of the effect of adrenomedullin on contraction of the urinary bladder due to acetylcholine.


[Figure 4] A diagram showing the amino acid sequence of adrenomedullin derived from human pheochromocytoma. RE1 to RE6 indicate fragments produced by digesting the amino acid sequence with arginylendopeptidase.

11-177549


-1-

[Name of the Document] DRAWINGS

[Fig. 1]



[Fig. 2]



11-177549

-2-


[Fig. 3]

* Contraction due to adding acetylcholine
is not significantly affected by the presence
of adrenomedullin.

Contraction due to acetylcholine
in the presence of 100nM of adrenomedullin

Contraction due to acetylcholine
↓
100nM~1μM

↓
100nM~1μM





Static tension of the urinary bladder
is not changed by adrenomedullin alone.

11-177549

-3-

[Fig. 4]





11-177549

- 1 -

[Name of the Document] ABSTRACT

[Abstract]

[Problems] Provision of a novel agent for promoting extension of bladder smooth muscle.

[Means for Solving the Problems] A composition for promoting extension of bladder smooth muscle, comprising adrenomedullin. The composition of the present invention may be used to ameliorate a urination disorder. The urination disorder may be a urinary incontinence selected from the group consisting of urge incontinence, reflex incontinence, and overflow incontinence.

[Selected Figure] None